

# Rangemate Custom 32-10 (R400) Clmnx

49C-3038

Type B Medicated Feed

42616

(For Beef Cattle Only)

For improved feed efficiency in growing-finishing cattle fed in confinement for slaughter.

### Active Drug Ingredient

Monensin (as Monensin sodium) 400.0 g/ton

### Guaranteed Analysis

Crude Protein		Min.	32.00 %
(This includes no more than 30.0 % equivalent protein from non-protein nitrogen)			
Crude Fat		Min.	10 %
Crude Fiber		Max.	0.1 %
Calcium (Ca)	Min.	0.05%	Max. 0.50 %
Phosphorus (P)		Min.	0.80 %
Salt	Min.	3.00%	Max. 4.00 %
Potassium (K)		Min.	1.50 %
Cobalt (Co)		Min.	4.6 ppm
Copper (Cu)		Min.	95 ppm
Iodine (I)		Min.	4.0 ppm
Manganese (Mn)		Min.	370 ppm
Zinc (Zn)		Min.	310 ppm
Selenium (Se)		Min.	2.0 ppm
Vitamin A		Min.	60,000 IU/lb
Vitamin D		Min.	15,000 IU/lb
Vitamin E		Min.	90 IU/lb
Dry Matter		Min.	55.0 %
Moisture		Max.	45.0 %

### Ingredients:

Molasses Products, Urea, Whey permeate, Corn Oil, Glycerin, Ammonium Polyphosphate, Magnesium chloride, Salt, Manganese polysaccharide complex, Zinc polysaccharide complex, Copper polysaccharide complex, Hydrolyzed yeast, Yeast extract, Yeast culture, Coconut oil, Maltodextrin, Lactic Acid, Xanthan gum, Sodium selenite, Cobalt Sulfate, Ethylenediamine Dihydriodide, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement.

**CANNOT BE FED FREE-CHOICE**

CAUTION: USE ONLY AS DIRECTED

Manufactured for Rangemate America LLC

701-667-8768

2475 26th St

Mandan, ND 585545

Liquid Bulk Shipment

## Mixing Directions

Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily, even when not in use.

### Directions for Use

Milligrams of Monensin per pound = 200.0

Minimum Feeding Rate - lbs/hd/day = 0.25  
Maximum Feeding Rate - lbs/hd/day = 2.40

### Caution

Inadequate mixing or agitation of monensin liquid Type B medicated feed has resulted in increased monensin concentration which has been fatal to cattle. Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Monensin medicated cattle feed is safe for use only in cattle. Consumption by unapproved species may result in toxic reaction. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

### Warning

Do not feed to lactating dairy cattle. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This product was manufactured in a facility that does not handle or store animal protein products prohibited in ruminant feed as described in 21 CFR section 589.2000.

Expiration Date: \_\_\_\_\_

8 weeks post manufacture